

# Strategic Initiatives for Flow Visualization in Environmental Sciences

Roxana Bujack<sup>1</sup> and Ariane Middel<sup>2</sup>

<sup>1</sup>Technical University Kaiserslautern, Germany

<sup>2</sup>Arizona State University, USA

## Abstract

*Flow plays a big role in the environmental sciences, but there are major differences between theoretically available and practically applied visualization techniques. This paper surveys various techniques in computational and environmental flow visualization, identifies challenges, and suggests strategic initiatives on how to bridge the gap.*

## 1. Introduction


Flow plays an important role in environmental sciences, because many of the Earth's physical and biological processes involve movement. In computer science, flow visualization is a well established research field that provides a variety of methods and algorithms that allow a human to see and understand the motion of water, air, or particles. Yet, a gap exists between the tools that have been proposed in flow visualization and those that have actually been applied to real world problems by domain experts in environmental sciences. In this paper, we investigate the differences between these two theoretically closely related fields, point out potential reasons for the gap, and suggest solutions to bridge it.

## 2. Flow Visualization

Flow visualization is the science of making flow fields visible [HJ05]. It is older than scientific computing [SL00, Mer12], but the use of computers has leveraged it into a new era, on which we will focus. Openly available programs, like ParaView [AGL\*05] or VisIt [CBW\*12], provide many flow visualization techniques. Even though there are overlaps, they can be structured into the following categories.

**Geometric Techniques.** Some of the most popular flow visualization techniques fall into the category of geometric techniques, Fig. 1. Here, geometric objects such as curves, surfaces, and volumes, are built from seeding and advecting massless particles to represent the behavior of the underlying flow [MLP\*09]. The basic curves that form the foundation of the geometric objects are stream, path, streak, and time lines [BS85]. Instead of visualizing the flow simply as lines, the curves can be colored and illuminated [ZSH96] and drawn in the shape of tubes or ribbons to encode additional information, e.g. rotation, divergence, or velocity [USM96]. Concatenation of the basic lines seeded along a starting curve results in stream surfaces [Hul92], path, time, or streak

surfaces [MLZ09, KGJ09]. Emphasis on divergence is well provided by flow polygons [SVL91] or flow volumes [MBC93], which copy experimental smoke advection.



(a) Ill. str. lines. (b) Str. tubes. (c) Str. ribbons. (d) Str. surfaces.

Figure 1: Different streamline-based geometric techniques applied to an example 3D flow field of swirling behavior.

**Texture-Based Techniques.** Dense and texture-based techniques are very popular for the visualization of 2D flow fields [LHD\*04, EGL\*06, NW13]. They cover the whole domain densely by transforming a texture along the flow and are inspired by the experimental flow visualization technique where oil is applied to parts in the wind tunnel to form schlieren. The first such method introduced was spot noise [VW91], where a set of spots are placed on the domain and smeared in flow direction, Fig. 2a. Line integral convolution (LIC) [CL93] is a similar technique, Fig. 2b. Instead of a finite set of spots, a dense texture is used as a starting point. The output texture is produced by a one-dimensional convolution of the input texture along the streamline passing through a pixel. Follow up research generalizes these fundamental techniques to better encode orientation [WG97], a variety of input textures [KB96], surfaces [FC95], 3D flow fields [IG97], time-varying flow data [SK97], and more efficient calculation [SH95]. Still, the main application remains to be 2D steady flow because of the induced clutter of a dense visualization technique.

**Topology-Based Techniques.** Vector field topology [HH89] separates the domain into areas in which all flow parcels have the




Figure 2: Different techniques applied to an example 2D flow field with a source, a saddle, and a sink.

same origin and destination, Fig. 2c. The topological skeleton consists of critical points (the positions that have zero velocity) and separatrices (their one-dimensional invariant manifolds) [PVH\*03, LHZP07]. The different flow patterns of vector field critical points can be categorized into saddles, sinks, sources, and vortices by the eigenvalues of their Jacobian [PC87]. It has become an important component of flow visualization, because it contains the important features of a vector field in a highly compressed representation [HH89]. Extensions have been made to the detection of higher order critical points [SHK\*97], separation and attachment lines [KHL99], closed streamlines [WS01], 3D vector fields [WTS\*05], and Galilean invariance [BHJ16]. It is also a means for flow field decomposition, simplification [TSH00], and design [The02]. Currently, the main challenge lies in generalizing flow topology to time varying data [PPF\*11].

**Feature-Based Techniques.** Most flow visualization techniques produce great results when applied to 2D fields, but fail in the 3D case due to clutter, because each drawn object occludes everything located behind it, Fig. 3. Feature based techniques reduce the amount of visualization elements to the important areas of a flow field [PVH\*03, EGL\*06]. This selection process can be imposed to all the techniques mentioned earlier. Specific detectors have been tailored to identify the most typical flow features, like vortices [Rot00], the elements of the flow topology [LHZP07], or shockwaves [WXWH13]. For more general purposes, vector field pattern detection algorithms [Ebl06, Sch11, BSH15, WSW16] allow the extraction of user defined features.

**Further Techniques.** There are other flow visualization techniques that can be considered as their own category, such as partition-




Figure 3: Feature based seeding of streamlines in regions of vortical behavior (b) vs equidistant seeding (a), (c).

based techniques [SJWS08], illustrative techniques [BCP\*12], visualization based on glyphs [PPVWS95, WPL96], Fig. 2d, or visualization of derived scalar quantities (e.g. velocity magnitude, vorticity, finite-time Ljapunov exponent, [HY00]).


### 3. Environmental Flow Visualization

Environmental science is, by nature, an interdisciplinary field, combining physical, biological, and information sciences to study environmental systems. Based on our literature review, we grouped environmental research into three categories: (1) atmospheric science; (2) ecology; and (3) geosciences. The following paragraphs review examples of applications in each category that involve time and space dependent movement data appropriate for flow visualization.


**Atmospheric Science.** Predominant atmospheric variables that pertain to flow are air and water, usually retrieved through CFD simulations. Windy [win16] is a web-based wind forecasting and visualization tool that maps wind direction and speed in Google Maps using a background image, color-coded by velocity, Fig. 4b. At a smaller spatial scale, [LPA11] conducted large-eddy simulations of wind farms and mapped simulated wind velocities as color-coded maps. They also provided contours of the time-averaged stream-wise velocity and turbulence intensity as color-coded isolines. [VHB10] illustrated microscale wind flow as color-coded contours of velocity magnitude in four horizontal planes, while [MHB\*14] displayed CFD modeling results for wind using arrows on a heat map, Fig. 4c. Another study investigated the impact of vertical wind shear on the predictability of tropical cyclones visualizing wind as arrows [ZT13]. Air quality assessment is another application area for flow visualization, as it is concerned with pollutant dispersion through wind. In [GZCL11], arrows are used as velocity vectors to display air-flow in urban street canyons. Modeling and analyzing wind flow is also important to assess the impact of hurricanes and tsunamis [PMD\*10, JS10, DBW\*10].

**Ecology.** The predominant application of flow visualization in ecology is animal migration. In a recent review, various tracking techniques for migratory birds were presented, ranging from GPS tracking to geologists [BTB\*11]. While bird tracking techniques were diverse, the presented migration visualizations were limited to a Google Earth overlay of migration paths and a 2D view of dotted lines on a radar image. Other applications used lines on a 2D map, color-coded by bird individuals [ESP\*10, SBvLP\*11].

**Geosciences.** Ocean flow simulation and analysis are crucial for



(a) Image courtesy of L. Treinish [Tre93].



(b) Wind forecasting in Windity [win16]. (c) ENVI-met [MHB\*14].


Figure 4: Wind direction and velocity mapping applications.

assessing environmental hazards, such as oil spills and sea trash. Previous research has predominantly used color-coding [MM98, Ain99, WPS\*16] and iso-lines [Sch02] to visualize ocean flow, Fig. 5. For coastal cities, flooding of urban areas was visualized using photo-realistic animations to explain impacts of sea level rise to decision-makers and the general public [GSH\*15]. A recent study presented Open Geosys, a tool to visualize geothermal energy and groundwater using streamlines and arrow glyphs [KBB\*12]. Flow visualization has also been used to illustrate glacier retreat [Kau12] and volcanoes [SYM14].


#### 4. Bridging the Gap

Our literature review suggests that environmental scientists generally use basic flow visualization techniques; atmospheric data are mapped as color-coded derived scalar fields or arrows, Fig. 4a and Fig. 4c, while ecologists prefer streamlines to visualize animal migration data. The gap between available flow visualization methods and their application seems to be smallest in geosciences, particularly ocean flows. In all three areas, higher visualization elements, like surfaces and volumes, feature based approaches, or topology are hardly used. We see the following reasons for this gap.

While fluid dynamics analysts often concentrate on relatively small, contained problems, environmental scientists deal with big, complex data sets from various sources in different formats. The data sets exhibit complex boundaries, are multidimensional, or stem from observations that are too sparse for higher visualization elements. For big data problems, environmental scientists often focus on the comparison of different parameters and variables, so the flow itself is visualized sparsely to share space with other variables (e.g., rain, snow, temperature) and topographic features (e.g., rivers, administrative boundaries). Often, environmental data sets are analyzed with respect to long term correlations, longitudinal trends, or



(a) Okubo Weiss.



(b) Kinetic energy.

Figure 5: Ocean visualization by color coding of derived scalar fields. Images courtesy of F. Samsel, M. Petersen, G. Abram with MPAS-Ocean, COSIM, LANL [WPS\*16].

averages; instantaneous flow patterns become secondary. Environmental scientists also have to address a broader, non-expert audience to communicate their findings, yielding more colorful, catchy, simplified, and sometimes animated visualizations and so far, we see fewer emphasis on animation and visual analytics in flow visualization [KMS\*08].

In our opinion, the demand for sparsity in comparative environmental visualization could be well satisfied by using feature based techniques and vector field topology. Environmental scientists would benefit from implementations that connect to their data formats and can easily be incorporated into existing tools the scientists depend on. Interaction might be difficult to achieve due to the size of the datasets, but GPU-based techniques, high performance computing, and scientific computing could offer solutions.

#### 5. Conclusion

We found that the two seemingly similar fields of flow visualization and environmental flow visualization are more different than expected. In environmental sciences, flow is almost always only visualized by basic techniques, like arrow glyphs, streamlines, or color coding of the velocity magnitude. This gap is mainly based on the demands of environmental sciences for sparse, catchy, and simplified visualizations that run fast on big datasets and are easy to integrate into existing environments. In the future it can be bridged if flow visualization investigates views in which the immediate flow may play a subordinate role and the environmental sciences consider feature-based and topological methods.

## Acknowledgements

This research was partly funded by the German Research Foundation (DFG) as part of the the IRTG 2057 "Physical Modeling for Virtual Manufacturing Systems and Processes".

## References

- [AGL\*05] AHRENS J., GEVECI B., LAW C., HANSEN C., JOHNSON C.: 36-paraview: An end-user tool for large-data visualization, 2005. 1
- [Ain99] AINSWORTH E.: Visualization of ocean colour and temperature from multi-spectral imagery captured by the japanese adeos satellite. *Journal of Visualization* 2, 2 (1999), 195–204. 3
- [BCP\*12] BRAMBILLA A., CARNECKY R., PEIKERT R., VIOLA I., HAUSER H.: Illustrative Flow Visualization: State of the Art, Trends and Challenges. *EG 2012 State of the Art Reports* (2012), 75–94. 2
- [BHJ16] BUJACK R., HLAWITSCHKA M., JOY K. I.: Topology-Inspired Galilean Invariant Vector Field Analysis. In *accepted for publication at IEEE Pacific Visualization Symposium, PacificVis 2016 in Taipei, Taiwan* (2016). 2
- [BHS15] BUJACK R., HOTZ I., SCHEUERMANN G., HITZER E.: Moment Invariants for 2D Flow Fields via Normalization in Detail. *IEEE Transactions on Visualization and Computer Graphics (TVCG)* 21, 8 (2015), 916–929. 2
- [BS85] BUNING P., STEGER J. L.: Graphics and flow visualization in computational fluid dynamics. In *Proceedings of the American Institute of Aeronautics and Astronautics 8th Computational Fluid Dynamics Conference* (1985), p. 814/820. 1
- [BTB\*11] BRIDGE E. S., THORUP K., BOWLIN M. S., CHILSON P. B., DIEHL R. H., FLÉRON R. W., HARTL P., KAYS R., KELLY J. F., ROBINSON W. D.: Technology on the move: recent and forthcoming innovations for tracking migratory birds. *BioScience* 61, 9 (2011), 689–698. 2
- [CBW\*12] CHILDS H., BRUGGER E., WHITLOCK B., MEREDITH J., AHERN S., PUGMIRE D., BIAGAS K., MILLER M., HARRISON C., WEBER G. H., KRISHNAN H., FOGAL T., SANDERSON A., GARTH C., BETHEL E. W., CAMP D., RÜBEL O., DURANT M., FAVRE J. M., NAVRÁTIL P.: VisIt: An End-User Tool For Visualizing and Analyzing Very Large Data. In *High Performance Visualization—Enabling Extreme-Scale Scientific Insight*. Oct 2012, pp. 357–372. 1
- [CL93] CABRAL B., LEEDOM L. C.: Imaging vector fields using line integral convolution. In *Proceedings of the 20th annual conference on Computer graphics and interactive techniques* (1993), SIGGRAPH '93, ACM, pp. 263–270. 1
- [DBW\*10] DIETRICH J. C., BUNYA S., WESTERINK J. J., EBERSOLE B. A., SMITH J. M., ATKINSON J. H., JENSEN R., RESIO D. T., LUETTICH R. A., DAWSON C.: A high-resolution coupled riverine flow, tide, wind, wind wave, and storm surge model for southern louisiana and mississippi. part II: Synoptic description and analysis of hurricanes katrina and rita. *Monthly Weather Review* 138, 2 (2010), 378–404. 2
- [Ebl06] EBLING J.: *Visualization and Analysis of Flow Fields using Clifford Convolution*. PhD thesis, University of Leipzig, Germany, 2006. 2
- [EGL\*06] ERLEBACHER G., GARTH C., LARAMEE R. S., THEISEL H., TRICOCHÉ X., WEINKAUF T., WEISKOPF D.: Texture and Feature-Based Flow Visualization - Methodology and Application. In *IEEE Visualization Tutorial* (2006). 1, 2
- [ESP\*10] EGEVANG C., STENHOUSE I. J., PHILLIPS R. A., PETERSEN A., FOX J. W., SILK J. R.: Tracking of arctic terns *Sterna paradisaea* reveals longest animal migration. *Proceedings of the National Academy of Sciences* 107, 5 (2010), 2078–2081. 2
- [FC95] FORSSELL L. K., COHEN S. D.: Using line integral convolution for flow visualization: Curvilinear grids, variable-speed animation, and unsteady flows. *Visualization and Computer Graphics, IEEE Transactions on* 1, 2 (1995), 133–141. 1
- [GSH\*15] GROTTTEL S., STAIB J., HEYER T., VETTER B., GUMHOLD S.: Real-Time Visualization of Urban Flood Simulation Data for Non-Professionals. In *Workshop on Visualisation in Environmental Sciences (EnvirVis)* (2015), Middel A., Rink K., Weber G. H., (Eds.), The Eurographics Association. 3
- [GZCL11] GU Z.-L., ZHANG Y.-W., CHENG Y., LEE S.-C.: Effect of uneven building layout on air flow and pollutant dispersion in non-uniform street canyons. *Building and Environment* 46, 12 (2011), 2657–2665. 2
- [HH89] HELMAN J., HESSELINK L.: Representation and display of vector field topology in fluid flow data sets. *Computer* 22, 8 (1989), 27–36. 1, 2
- [HJ05] HANSEN C., JOHNSON C.: *The Visualization Handbook*. Referex Engineering. Butterworth-Heinemann, 2005. 1
- [Hul92] HULTQUIST J. P.: Constructing stream surfaces in steady 3d vector fields. In *Proceedings of the 3rd conference on Visualization'92* (1992), IEEE Computer Society Press, pp. 171–178. 1
- [HY00] HALLER G., YUAN G.: Lagrangian coherent structures and mixing in two-dimensional turbulence. *Phys. D* 147, 3-4 (2000), 352–370. 2
- [IG97] INTERRANTE V., GROSCH C.: Strategies for effectively visualizing 3d flow with volume lic. In *Proceedings of the 8th conference on Visualization'97* (1997), IEEE Computer Society Press, pp. 421–ff. 1
- [JS10] JAIMES B., SHAY L. K.: Near-inertial wave wake of hurricanes katrina and rita over mesoscale oceanic eddies. *Journal of Physical Oceanography* 40, 6 (2010), 1320–1337. 2
- [Kau12] KAUFMANN V.: The evolution of rock glacier monitoring using terrestrial photogrammetry: the example of äusseres hochebenkar rock glacier (austria). *Austrian Journal of Earth Sciences* 105, 2 (2012), 63–77. 3
- [KB96] KIU M.-H., BANKS D. C.: Multi-frequency noise for lic. In *Proceedings of the 7th Conference on Visualization'96* (1996), IEEE Computer Society Press, pp. 121–126. 1
- [KBB\*12] KOLDITZ O., BAUER S., BILKE L., BÖTTCHER N., DELFS J.-O., FISCHER T., GÖRKE U. J., KALBACHER T., KOSAKOWSKI G., MCDERMOTT C., ET AL.: Opengeosys: an open-source initiative for numerical simulation of thermo-hydro-mechanical/chemical (thm/c) processes in porous media. *Environmental Earth Sciences* 67, 2 (2012), 589–599. 3
- [KGJ09] KRISHNAN H., GARTH C., JOY K.: Time and Streak Surfaces for Flow Visualization in Large Time-Varying Data Sets. *IEEE Transactions on Visualization and Computer Graphics* 15, 6 (2009), 1267–1274. 1
- [KHL99] KENWRIGHT D., HENZE C., LEVIT C.: Feature Extraction of Separation and Attachment Lines. *Visualization and Computer Graphics, IEEE Transactions on* 5, 2 (1999), 135–144. 2
- [KMS\*08] KEIM D. A., MANSMANN F., SCHNEIDEWIND J., THOMAS J., ZIEGLER H.: *Visual analytics: Scope and challenges*. Springer, 2008. 3
- [LHD\*04] LARAMEE R. S., HAUSER H., DOLEISCH H., VROLIJK B., POST F. H., WEISKOPF D.: The State of the Art in Flow Visualization: Dense and Texture-Based Techniques. *Computer Graphics Forum* 23 (2004), 2004. 1
- [LHZP07] LARAMEE R. S., HAUSER H., ZHAO L., POST F. H.: Topology-Based Flow Visualization, The State of the Art. In *Topology-based Methods in Visualization* (2007), pp. 1–19. 2
- [LPA11] LU H., PORTÉ-AGEL F.: Large-eddy simulation of a very large wind farm in a stable atmospheric boundary layer. *Physics of Fluids (1994-present)* 23, 6 (2011), 065101. 2
- [MBC93] MAX N., BECKER B., CRAWFIS R.: Flow volumes for interactive vector field visualization. In *Proceedings of the 4th conference on Visualization'93* (1993), IEEE Computer Society, pp. 19–24. 1
- [Mer12] MERZKIRCH W.: *Flow visualization*. Elsevier, 2012. 1

- [MHB\*14] MIDDEL A., HÄB K., BRAZEL A. J., MARTIN C. A., GUHATHAKURTA S.: Impact of urban form and design on mid-afternoon microclimate in phoenix local climate zones. *Landscape and Urban Planning* 122 (2014), 16–28. 2, 3
- [MLP\*09] MCLOUGHLIN T., LARAMEE R. S., PEIKERT R., POST F. H., CHEN M.: Over Two Decades of Integration-Based, Geometric Flow Visualization. In *EG 2009 - State of the Art Reports* (2009), pp. 73–92. 1
- [MLZ09] MCLOUGHLIN T., LARAMEE R. S., ZHANG E.: Easy integral surfaces: a fast, quad-based stream and path surface algorithm. In *Proceedings of the 2009 Computer Graphics International Conference* (2009), ACM, pp. 73–82. 1
- [MM98] MCPHERSON A., MALTRUD M.: Poptex: Interactive ocean model visualization using texture mapping hardware. In *Proceedings of the conference on Visualization '98* (1998), IEEE Computer Society Press, pp. 471–474. 3
- [NW13] NETZEL R., WEISKOPF D.: Texture-Based Flow Visualization. *Computing in Science and Engineering* 15, 6 (2013), 96–102. 1
- [PC87] PERRY A., CHONG M.: A description of eddying motions and flow patterns using critical-point concepts. *Annual Review of Fluid Mechanics* 19, 1 (1987), 125–155. 2
- [PMD\*10] POWELL M. D., MURILLO S., DODGE P., UHLHORN E., GAMACHE J., CARDONE V., COX A., OTERO S., CARRASCO N., AN-NANE B.: Reconstruction of hurricane katrina's wind fields for storm surge and wave hindcasting. *Ocean Engineering* 37, 1 (2010), 26–36. 2
- [PPF\*11] POBITZER A., PEIKERT R., FUCHS R., SCHINDLER B., KUHN A., THEISEL H., MATKOVIC K., HAUSER H.: The State of the Art in Topology-based Visualization of Unsteady Flow. *Computer Graphics Forum* 30, 6 (2011), 1789–1811. 2
- [PPVWS95] POST F. H., POST F. J., VAN WALSUM T., SILVER D.: Iconic techniques for feature visualization. In *Proceedings of the 6th conference on Visualization '95* (1995), IEEE Computer Society, p. 288. 2
- [PVH\*03] POST F. H., VROLIJK B., HAUSER H., LARAMEE R. S., DOLEISCH H.: The State of the Art in Flow Visualisation: Feature Extraction and Tracking. *Computer Graphics Forum* 22, 4 (2003), 775–792. 2
- [Rot00] ROTH M.: *Automatic Extraction of Vortex Core Lines and Other Line-Type Features for Scientific Visualization*. PhD Dissertation No. 13673, ETH Zurich, 2000. published by Hartung-Gorre Verlag, Konstanz, ISBN 3-89649-582-8. 2
- [SBvLP\*11] SHAMOUN-BARANES J., VAN LOON E. E., PURVES R. S., SPECKMANN B., WEISKOPF D., CAMPHUYSEN C. J.: Analysis and visualization of animal movement. *Biology letters* (2011). 2
- [Sch02] SCHLITZER R.: Interactive analysis and visualization of geoscience data with ocean data view. *Computers & geosciences* 28, 10 (2002), 1211–1218. 3
- [Sch11] SCHLEMMER M.: *Pattern Recognition for Feature Based and Comparative Visualization*. PhD thesis, Universität Kaiserslautern, Germany, 2011. 2
- [SH95] STALLING D., HEGE H.-C.: Fast and resolution independent line integral convolution. In *Proceedings of the 22nd annual conference on Computer graphics and interactive techniques* (1995), ACM, pp. 249–256. 1
- [SHK\*97] SCHEUERMANN G., HAGEN H., KRÜGER H., MENZEL M., ROCKWOOD A.: Visualization of higher order singularities in vector fields. In *Proceedings of the 8th conference on Visualization '97* (1997), IEEE Computer Society Press, pp. 67–74. 2
- [SJWS08] SALZBRUNN T., JÄNICKE H., WISCHGOLL T., SCHEUERMANN G.: The State of the Art in Flow Visualization: Partition-based Techniques. In *Simulation and Visualization 2008 Proceedings* (2008). 2
- [SK97] SHEN H.-W., KAO D. L.: Uffic: a line integral convolution algorithm for visualizing unsteady flows. In *Proc. of the 8th conference on Visualization '97* (1997), IEEE Computer Society Press, pp. 317–ff. 1
- [SL00] SMITS A. J., LIM T. T.: *Flow Visualization Techniques and Examples*. Imperil College Press, London, UK, 2000. 1
- [SVL91] SCHROEDER W. J., VOLPE C. R., LORENSEN W. E.: The stream polygon: A technique for 3d vector field visualization. In *Proceedings of the 2nd conference on Visualization '91* (1991), IEEE Computer Society Press, pp. 126–132. 1
- [SYM14] SAUER F., YU H., MA K.-L.: Trajectory-based flow feature tracking in joint particle/volume datasets. *Visualization and Computer Graphics, IEEE Transactions on* 20, 12 (2014), 2565–2574. 3
- [The02] THEISEL H.: Designing 2d vector fields of arbitrary topology. In *Computer Graphics Forum* (2002), vol. 21, Wiley Online Library, pp. 595–604. 2
- [Tre93] TREINISH L. A.: Visualization of stratospheric ozone depletion and the polar vortex. In *Visualization, 1993. Visualization '93, Proceedings., IEEE Conference on* (1993), IEEE, pp. 391–396. 3
- [TSH00] TRICOCHÉ X., SCHEUERMANN G., HAGEN H.: A topology simplification method for 2d vector fields. In *Visualization 2000. Proceedings* (2000), IEEE, pp. 359–366. 2
- [USM96] UENG S. K., SIKORSKI C., MA K. L.: Efficient streamline, streamribbon, and streamtube constructions on unstructured grids. *Visualization and Computer Graphics, IEEE Transactions on* 2, 2 (1996), 100–110. 1
- [VHB10] VAN HOOFF T., BLOCKEN B.: Coupled urban wind flow and indoor natural ventilation modelling on a high-resolution grid: A case study for the amsterdam ArenA stadium. *Environmental Modelling & Software* 25, 1 (2010), 51–65. 2
- [VW91] VAN WIJK J. J.: Spot noise texture synthesis for data visualization. In *ACM Siggraph Computer Graphics* (1991), vol. 25, ACM, pp. 309–318. 1
- [WG97] WEGENKITTL R., GRÖLLER E.: Fast oriented line integral convolution for vector field visualization via the internet. In *Proceedings of the 8th conference on Visualization '97* (1997), IEEE Computer Society Press, pp. 309–316. 1
- [win16] Windy, SE, wind map & forecast, 2016. URL: <https://www.windyty.com>. 2, 3
- [WPL96] WITTENBRINK C. M., PANG A. T., LODHA S. K.: Glyphs for visualizing uncertainty in vector fields. *Visualization and Computer Graphics, IEEE Transactions on* 2, 3 (1996), 266–279. 2
- [WPS\*16] WOODRING J., PETERSEN M., SCHMEISSER A., PATCHETT J., AHRENS J., HAGEN H.: In situ eddy analysis in a high-resolution ocean climate model. *Visualization and Computer Graphics, IEEE Transactions on* 22, 1 (2016), 857–866. 3
- [WS01] WISCHGOLL T., SCHEUERMANN G.: Detection and visualization of closed streamlines in planar flows. *Visualization and Computer Graphics, IEEE Transactions on* 7, 2 (2001), 165–172. 2
- [WSW16] WANG Z., SEIDEL H.-P., WEINKAUF T.: Multi-field pattern matching based on sparse feature sampling. *Visualization and Computer Graphics, IEEE Transactions on* 22, 1 (2016), 807–816. 2
- [WTS\*05] WEINKAUF T., THEISEL H., SHI K., HEGE H.-C., SEIDEL H.-P.: Extracting Higher Order Critical Points and Topological Simplification of 3D Vector Fields. In *Proc. IEEE Visualization 2005* (Minneapolis, U.S.A., 2005), pp. 559–566. 2
- [WXWH13] WU Z., XU Y., WANG W., HU R.: Review of Shock Wave Detection Method in {CFD} Post-Processing. *Chinese Journal of Aeronautics* 26, 3 (2013), 501 – 513. 2
- [ZSH96] ZÖCKLER M., STALLING D., HEGE H.-C.: Interactive visualization of 3d-vector fields using illuminated stream lines. In *Visualization '96. Proceedings.* (1996), IEEE, pp. 107–113. 1
- [ZT13] ZHANG F., TAO D.: Effects of vertical wind shear on the predictability of tropical cyclones. *Journal of the Atmospheric Sciences* 70, 3 (2013), 975–983. 2